

Adolescent Vaccine Schedule 11-18 Years Old



Vaccine	Dose	Timing of Immunization
Hepatitis B (Hep B)¹	3 doses	1st dose / 2nd dose 1 month after 1st dose 3rd dose 4-6 months after 2nd dose
Measles, Mumps, Rubella (MMR)²	2 doses	1st dose / 2nd dose 1 month after 1st dose
Varicella (Chickenpox)³	2 doses	1st dose/ 2nd dose 1-3 months after 1st dose depending on age
Hepatitis A (Hep A)⁴	2 doses	1st dose/ 2nd dose 6-12 months after 1st dose
Influenza (Flu)⁵	1 dose	1 dose yearly
Pneumococcal⁶	1 dose	1 dose to adolescents with serious diseases
Meningococcal Conjugate (MenACWY)⁷	2 doses	1 dose is recommended at 11-12 years of age with booster dose at 16 years of age
Tdap (Tetanus, Diphtheria, Pertussis)⁸	1 dose	1 dose of Tdap is recommended at 11-12 years of age
Human Papillomavirus (HPV)⁹	2 doses or 3 doses	Age 9 through 14 years at initial vaccination: 1st dose / 2nd dose 6-12 months after 1st dose Age 15 years or older at initial vaccination: 1st dose / 2nd dose 1-2 months after 1st dose 3rd dose 6 months after 1st dose
Meningococcal B¹⁰	2 doses or 3 doses	1st dose/ 2nd dose 1 month after 1st dose 1st dose / 2nd dose 6 months after 1st dose 3 doses 4 months after 2nd dose
COVID-19¹¹	1 dose	1 dose yearly

See reverse side for detailed recommendations

-
- 1 Hepatitis B:** Three doses of the hepatitis B vaccine are needed for all adolescents. A two-dose hepatitis B vaccine is available for adolescents 11-15 years of age.
 - 2 Measles, Mumps, Rubella (MMR):** Two doses of the MMR vaccine are recommended to provide full protection for adolescents who have not previously been vaccinated. The MMR vaccine should not be given to adolescents who are known to be pregnant or who are considering becoming pregnant within one month of vaccination.
 - 3 Varicella (Chickenpox):** The chickenpox vaccine is recommended for anyone who has not had chickenpox disease or has not had the chickenpox vaccine. The chickenpox vaccine should not be given to adolescents who are known to be pregnant or who are considering becoming pregnant within one month of vaccination.
 - 4 Hepatitis A:** Two doses of the hepatitis A vaccine are recommended for those not previously vaccinated.
 - 5 Influenza (Flu):** A yearly flu vaccine is recommended for all teens.
 - 6 Pneumococcal Vaccine:** A single dose of the pneumococcal conjugate vaccine is recommended for teens with serious diseases such as diabetes, leukemia, heart disease, lung disease, HIV infection, or AIDS.
 - 7 Meningococcal (Meningitis):** Two doses of the Meningococcal Conjugate Vaccine (MCV) are recommended for adolescents. The first dose is recommended at 11-12 years of age with a booster at age 16. Teens who received MCV for the first time at 13 through 15 years of age will need a one-time booster between 16 and 18 years of age.
 - 8 Tdap (Tetanus, Diphtheria, Pertussis):** One dose of the Tdap vaccine is recommended for teens. Adolescents 11-18 years of age who have not received Tdap vaccine should receive a single dose of Tdap. One dose of Tdap vaccine is recommended during each pregnancy (preferred during 27 through 36 weeks gestation), regardless of time since prior Td or Tdap vaccination.
 - 9 Human Papillomavirus (HPV):** Depending on the age at initial vaccination, two or three doses of the HPV vaccine are recommended for all adolescents at 11-12 years of age. Catch-up vaccination is recommended for those aged 13-18 years of age who were not previously vaccinated or have not completed the full vaccine series.
 - 10 Meningococcal B:** Based on clinical shared decision-making, 2 or 3 (depending on the brand) doses of the Meningococcal B vaccine are recommended for adolescents aged 16-23 (preferably 16-18) years, who are at increased risk for meningococcal disease.
 - 11 COVID-19:** A yearly COVID-19 vaccine is recommended for all teens.